

Medication Guide

BRIXADI™ (brix-a'-dee) (buprenorphine) extended-release injection, for subcutaneous use (CIII)

What is the most important information I should know about BRIXADI?

- BRIXADI contains a medicine called buprenorphine. Buprenorphine is an opioid that can cause serious and life-threatening problems, especially if you take or use certain other medicines or drugs.
- Talk to your healthcare provider about naloxone. Naloxone is a medicine that is available to patients for the emergency treatment of an opioid overdose. If naloxone is given, you must call 911 or get emergency medical help right away to treat an overdose or accidental use of an opioid.
- Because of the serious risk of potential harm or death from self-injecting BRIXADI into a vein (intravenously), it is only available through a controlled program called the BRIXADI REMS Program.
 - BRIXADI is not available in retail pharmacies.
 - Your BRIXADI injection will only be given to you by a healthcare provider.
- **BRIXADI can cause serious and life-threatening problems. Get emergency help right away if you:**
 - feel faint
 - feel dizzy
 - are confused
 - feel sleepy or uncoordinated
 - have blurred vision
 - have slurred speech
 - are breathing slower than normal
 - cannot think well or clearly
- **Do not take BRIXADI with certain medicines. Taking BRIXADI with other medicines, benzodiazepines, alcohol, other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.** In an emergency, have family members tell the emergency department staff that you are physically dependent on an opioid and are being treated with BRIXADI.

What is BRIXADI?

BRIXADI is a prescription medicine used to treat moderate to severe addiction in people who:

- have started treatment with a single dose of a buprenorphine medicine in the form of a sublingual tablet or buccal film (transmucosal), OR
- who are already being treated with buprenorphine

BRIXADI should be used as part of a complete treatment plan that also includes counseling and behavioral therapy.

Who should not use BRIXADI?

Do not use BRIXADI if you are allergic to buprenorphine or any ingredients in BRIXADI.

Caution: The packaging of this product contains natural rubber latex which may cause allergic reactions.

Before receiving BRIXADI, tell your healthcare provider about all of your medical conditions, including if you have:

- trouble breathing or lung problems
- a curve in your spine that affects your breathing
- Addison's disease
- an enlarged prostate (men)
- problems urinating
- liver, kidney, or gallbladder problems
- a history of alcoholism
- a head injury or brain problem
- mental health problems
- adrenal gland or thyroid gland problems

Tell your healthcare provider if you are:

- **pregnant or plan to become pregnant.** If you use BRIXADI while pregnant, your baby may have symptoms of opioid withdrawal at birth that could be life-threatening if not recognized and treated.
- **breastfeeding.** BRIXADI can pass into your breast milk and may harm your baby. Talk to your healthcare provider about the best way to feed your baby if you take BRIXADI. Monitor your baby for increased drowsiness and breathing problems if you breastfeed during treatment with BRIXADI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

You may have detectable levels of BRIXADI in your body for several months after stopping treatment with BRIXADI.

How will I receive BRIXADI?

- You will receive BRIXADI by your healthcare provider as an injection just under the skin (subcutaneous) of your buttock, thigh, stomach (abdomen), or upper arm. If you are new to buprenorphine treatment, the upper arm should only be used after several doses of BRIXADI.
- You will receive BRIXADI 1 time every week or 1 time every month.
- BRIXADI is injected as a liquid. After the injection, BRIXADI changes to a gel form called a depot. The depot is not always felt under the skin.
- Do not try to remove the depot.
- If you miss a dose of BRIXADI, see your healthcare provider to get your BRIXADI injection as soon as possible.

What should I avoid while receiving BRIXADI?

- **Do not drive, operate heavy machinery or perform any other dangerous activities until you know how BRIXADI affects you.** BRIXADI can cause drowsiness and slow reaction times. BRIXADI can make you sleepy, dizzy, or lightheaded. This may happen more often in the first day after your injection and when your dose is being changed.
- **You should not drink alcohol or use prescription or over-the-counter medicines that contain alcohol** during treatment with BRIXADI, because this can lead to loss of consciousness or even death.

What are the possible side effects of BRIXADI?

BRIXADI can cause serious side effects, including:

- **Trouble breathing.** Taking BRIXADI with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants can cause breathing problems that can lead to coma and death.
- **Sleepiness, dizziness, and problems with coordination.**
- **Physical dependence or abuse.**
- **Liver problems.** Call your healthcare provider right away if you notice any of these symptoms:

- your skin or the white part of your eyes turns yellow
- dark or “tea colored” urine
- light colored stools (bowel movements)
- loss of appetite
- pain, aching, or tenderness on the right side of your stomach area
- nausea

Your healthcare provider should do tests to check the liver before and during treatment with BRIXADI.

- **Allergic reaction.** You may have a rash, hives, swelling of your face, wheezing, light-headedness when changing positions, feeling faint, or loss of consciousness. Call your healthcare provider or get emergency help right away.
- **Opioid withdrawal.** Call your healthcare provider right away if you get any of these symptoms:
 - shaking
 - sweating more than normal
 - feeling hot or cold more than normal
 - runny nose
 - watery eyes
 - goose bumps
 - diarrhea
 - vomiting
 - muscle aches

These symptoms may start weeks to months after your last dose of BRIXADI. Tell your healthcare provider if you develop any of these symptoms.

- **Decrease in blood pressure.** You may feel dizzy if you get up too fast from sitting or lying down.

The most common side effects of BRIXADI include:

- injection site pain
- headache
- constipation
- nausea
- injection site itching
- injection site redness
- trouble sleeping (insomnia)
- urinary tract infection

BRIXADI may affect fertility in males and females. Talk to your healthcare provider if this is a concern for you.

These are not all the of possible side effects of BRIXADI.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of BRIXADI.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use BRIXADI for a condition for which it was not prescribed. You can ask your pharmacist or healthcare provider for information that is written for healthcare professionals.

Distributed by Braeburn, Inc., Plymouth Meeting, PA 19462, USA.
BRIX-002-01 Based on PI BRX-001-01 **XX/21**
© 2020 Braeburn, Inc. BRIXADI™ is a trademark of Braeburn Inc.
For more information, go to [www. BRIXADI.com](http://www.BRIXADI.com) or call 1-833-274-9234.